

Discrete $\mathcal{PT}$ –symmetric square-well oscillators

Miloslav Znojil¹

Ústav jaderné fyziky AV ČR, 250 68 Řež, Czech Republic

Abstract

Exact solvability of the discretized N –point version of the $\mathcal{PT}$ –symmetric square-well model at all N is pointed out. Its wave functions are found proportional to the classical Tshebyshev polynomials $U_k \cos \theta$ of a complex argument. A compact secular equation is derived giving the real spectrum of energies at all the non-Hermiticity strengths $Z \in (-Z_{crit}(N), Z_{crit}(N))$. In the limit $Z \rightarrow 0$ the model degenerates to a Hermitian Hückel Hamiltonian.

PACS 03.65.Ge, 02.60.Lj, 02.70.Bf, 31.15.Ct

¹e-mail: znojil@ujf.cas.cz

1 Introduction

The essence of the concept of the so called $\mathcal{PT}$ –symmetric Quantum Mechanics [1] lies in a half-forgotten fact that although the operators of observables (say, Hamiltonians H) *must* be Hermitian (with respect to a metric Θ in Hilbert space), they are *also* allowed to be Hermitian with respect to a *nontrivial* metric,

$$H^\dagger = \Theta H \Theta^{-1}, \quad I \neq \Theta = \Theta^\dagger > 0. \quad (1)$$

In sufficient detail, the formal aspects of this idea were already well described in the review paper [2] showing that in nuclear physics a decisive simplification of certain Schrödinger equations may be achieved after a formal transition from the most common $\Theta_1 = I$ to another $\Theta_2 \neq I$.

Beyond the area of nuclear physics the feasibility of the necessary calculations represents a key technical challenge for $\Theta \neq I$. Fortunately, for some differential Hamiltonians $H = \hat{p}^2 + V(x)$, an unexpectedly satisfactory answer has been found in a factorization of Θ , typically, into a product of parity $\mathcal{P}$ and the so called quasi-parity $\mathcal{Q}$ [3] or charge $\mathcal{C}$ [4]. In the other words, one requires that the observables are *both* $\mathcal{PT}$ –symmetric and $\mathcal{CP}$ –pseudo-Hermitian. This means that all our non-Hermitian observables must be compatible with eq. (1) *and* that they must also commute with certain operator $\mathcal{PT}$. In the terminology advocated by A. Mostafazadeh [5] the latter requirement should be generalized and re-interpreted as the so called $\mathcal{P}$ –pseudo-Hermiticity relation

$$H^\dagger = \mathcal{P} H \mathcal{P}^{-1}, \quad I \neq \mathcal{P} = \mathcal{P}^\dagger \quad (2)$$

In such a widely accepted scenario [6] our observables may remain manifestly non-Hermitian in the current sense, $H \neq H^\dagger$. Still, their spectra *must* remain real and the work with the underlying indefinite ‘pseudo-metric’ operators of generalized parity $\mathcal{P}$ *should* remain sufficiently easy.

In the light of these two requirements we feel particularly inspired here

- [a] by the rigorous Krein-space analysis [7] of the reality of spectra of quantum particles confined inside a one-dimensional $\mathcal{PT}$ –symmetric box $V(x)$,

- **[b]** in context **[a]**, by our older explicit construction [8] of bound states in one of the most elementary square-well forms of $V(x)$,
- **[c]** in context **[b]**, by the very recent Weigert's [9] three-by-three matrix discretization

$$\begin{pmatrix} 2 + \frac{1}{4}iZ & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 - \frac{1}{4}iZ \end{pmatrix} \begin{pmatrix} \alpha_0 \\ \gamma \\ \beta_0 \end{pmatrix} = \frac{1}{4}E \begin{pmatrix} \alpha_0 \\ \gamma \\ \beta_0 \end{pmatrix}. \quad (3)$$

of the oscillator of ref. [8].

In this general framework we intend to start from the $\mathcal{PT}$ -symmetric ordinary differential Schrödinger equation

$$\left[-\frac{d^2}{dx^2} + V(x) \right] \psi(x) = E \psi(x), \quad \psi(\pm 1) = 0, \quad V(x) = [V(-x)]^* \quad (4)$$

and from the proof given in ref. [7] that the resulting spectrum of energies $E = E_n$, $n = 0, 1, \dots$ is *real* and discrete for *all* the complex $\mathcal{PT}$ -symmetric potentials $V(x)$ which are not too strong,

$$\|V\|_\infty < \frac{3}{8}\pi^2 \approx 3.701. \quad (5)$$

In the spirit of item **[b]** we shall only pay attention to one of the simplest piecewise constant and purely imaginary antisymmetric potentials

$$V(x) = \begin{cases} +iZ & x \in (-1, 0), \\ -iZ & x \in (0, 1). \end{cases} \quad (6)$$

Along the lines of item **[c]** we shall introduce the Runge-Kutta discrete lattice of coordinates,

$$x_0 = -1, \quad x_k = x_{k-1} + h = -1 + kh, \quad h = \frac{2}{N}, \quad k = 1, 2, \dots, N$$

and pay attention to the general discrete analogue

$$-\frac{\psi(x_{k+1}) - 2\psi(x_k) + \psi(x_{k-1}))}{h^2} - i \operatorname{sign}(x_k)Z \psi(x_k) = E \psi(x_k) \quad (7)$$

of our differential Schrödinger equation (4) + (6). In combination with the boundary conditions

$$\psi(x_0) = \psi(x_N) = 0,$$

this in fact represents a generalization of the Weigert's $N = 4$ eq. (3) to all the integers N .

2 Models with the even $N = 2n + 4$

Let us recollect that the continuous $N = \infty$ model (6) is exactly solvable [8]. The solvability in closed form also characterizes its modifications with periodic boundary conditions and/or more discontinuities [10]. In this context, the exact solvability of the following $(N - 1)$ -dimensional matrix generalization

$$\left(\begin{array}{ccc|c|ccc} i\xi - F & -1 & & & & & \\ & -1 & i\xi - F & \ddots & & & \\ & & \ddots & \ddots & -1 & & \\ & & & -1 & i\xi - F & -1 & \\ \hline & & & -1 & -F & -1 & \\ \hline & & & -1 & -i\xi - F & \ddots & \\ & & & & \ddots & \ddots & -1 \\ & & & & & -1 & -i\xi - F \end{array} \right) \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \vdots \\ \alpha_n \\ \gamma \\ \beta_n \\ \vdots \\ \beta_0 \end{pmatrix} = 0 \quad (8)$$

of the three-dimensional Weigert's model (3) with the re-scaled energy eigenvalues $F = E h^2 - 2$ and with the re-scaled strength $\xi = Z h^2$ of the non-Hermiticity would not be too surprising.

In the first step of our analysis, due to the $\mathcal{PT}$ -symmetry of our problem we may set

$$\alpha_k = a_k + i b_k, \quad \beta_k = a_k - i b_k \equiv \alpha_k^*, \quad k = 0, 1, \dots, n$$

with some *real* elements $\gamma = \psi(0)$, $a_k = \text{Re } \psi(x_{k+1})$ and $b_k = \text{Im } \psi(x_{k+1})$. Once we recollect the definition of the classical Tshebyshev polynomials of the second kind

[11],

$$U_k(\cos \theta) = \frac{\sin(k+1)\theta}{\sin \theta}, \quad k = 0, 1, \dots,$$

we immediately obtain the wave functions in closed form,

$$\alpha_k = U_k \left(\frac{-F + i\xi}{2} \right) (a + ib), \quad k = 0, 1, \dots, n. \quad (9)$$

This reduces the full tridiagonal $(N-1) \times (N-1)$ -dimensional matrix eq. (8) to the mere three matching conditions,

$$\begin{pmatrix} -1 & i\xi - F & -1 & 0 & 0 \\ 0 & -1 & -F & -1 & 0 \\ 0 & 0 & -1 & -i\xi - F & -1 \end{pmatrix} \begin{bmatrix} U_{n-1} \left(\frac{-F+i\xi}{2} \right) (a + ib) \\ U_n \left(\frac{-F+i\xi}{2} \right) (a + ib) \\ \gamma \\ U_n \left(\frac{-F-i\xi}{2} \right) (a - ib) \\ U_{n-1} \left(\frac{-F-i\xi}{2} \right) (a - ib) \end{bmatrix} = 0. \quad (10)$$

The first and the third lines may be simplified to give

$$\gamma = U_{n+1} \left(\frac{-F + i\xi}{2} \right) (a + ib) = U_{n+1} \left(\frac{-F - i\xi}{2} \right) (a - ib). \quad (11)$$

The middle line defines the product

$$F \gamma = -U_n \left(\frac{-F + i\xi}{2} \right) (a + ib) - U_n \left(\frac{-F - i\xi}{2} \right) (a - ib). \quad (12)$$

This forces us to separate the $F = 0$ case as exceptional.

2.1 The existence of a nontrivial solution at $F = 0$

Once we set, tentatively, $F = 0$, it is easy to deduce from eq. (11) that the parameter a must vanish for the even $n = 0, 2, 4, \dots$ (and we may normalize $b = 1$) while $b = 0$ and $a = 1$ for the odd $n = 1, 3, 5, \dots$. Thus, eq. (11) degenerates to the mere definition of the last element γ of the eigenvector and we are left with the single secular eq. (12) which acquires the following two alternative forms,

$$\begin{aligned} U_n \left(\frac{1}{2} i\xi \right) - U_n \left(\frac{1}{2} i\xi \right) &= 0, & n = 2m, \\ U_n \left(\frac{1}{2} i\xi \right) + U_n \left(-\frac{1}{2} i\xi \right) &= 0, & n = 2m + 1. \end{aligned} \quad (13)$$

At any $m = 0, 1, \dots$ these conditions are satisfied identically. We may conclude that our tentative “guess of the energy” was correct and that $F = 0$ is always the eigenvalue. It is remarkable that in spite of the non-Hermiticity of the Hamiltonian, this “robust” eigenvalue remains real at *all* the real couplings $Z \in (-\infty, \infty)$.

3 Closed secular equations for $N = 2n + 4$

Whenever $F \neq 0$ we may treat eq. (11) not only as the condition of vanishing of the imaginary part of γ ,

$$U_{n+1} \left(\frac{-F + i\xi}{2} \right) (a + ib) = U_{n+1} \left(\frac{-F - i\xi}{2} \right) (a - ib) \quad (14)$$

but also as an explicit definition of the non-vanishing left-hand-side quantity $F\gamma$ in eq. (12). Its insertion simplifies the latter relation,

$$T_{n+1} \left(\frac{-F + i\xi}{2} \right) (a + ib) = -T_{n+1} \left(\frac{-F - i\xi}{2} \right) (a - ib) \quad (15)$$

where $T_k(z)$ denotes the k -th Tshebyshev polynomial of the first kind.

One of the latter two relations defines the normalization vector $(a, b) = (a_0, b_0)$ while their ratio gives

$$T_{n+1} \left(\frac{-F + i\xi}{2} \right) U_{n+1} \left(\frac{-F - i\xi}{2} \right) + T_{n+1} \left(\frac{-F - i\xi}{2} \right) U_{n+1} \left(\frac{-F + i\xi}{2} \right) = 0. \quad (16)$$

This is our final secular equation which defines, in an implicit manner, the energies F as functions of the couplings ξ .

An efficient numerical treatment of the latter eigenvalue problem may be based on the re-parametrization

$$\frac{-F + i\xi}{2} = \cos \varphi, \quad \operatorname{Re} \varphi = \alpha, \quad \operatorname{Im} \varphi = \beta \quad (17)$$

i.e.,

$$\frac{1}{2} F = -\cos \alpha \cosh \beta, \quad \frac{1}{2} \xi = -\sin \alpha \sinh \beta. \quad (18)$$

In opposite direction, the inversion of this change of variables

$$\cos \alpha = -\frac{1}{2 \cosh \beta} F, \quad \sinh \beta = \frac{1}{2\sqrt{2}} \sqrt{F^2 + \xi^2 - 4 + \sqrt{(F^2 + \xi^2 - 4)^2 + 16 \xi^2}}.$$

transforms eq. (16) into the compact and transparent trigonometric secular equation

$$\operatorname{Re} \frac{\sin[(n+1)\varphi] \cos[(n+1)\varphi^*]}{\sin \varphi} = 0. \quad (19)$$

Its roots may be determined, numerically, as lying in the domain with negative $\beta < 0$ and with $\alpha \in (0, \pi/2)$ for the negative $F < 0$ and with $\alpha \in (\pi/2, \pi)$ for the positive $F > 0$. In this picture, the constant value of the coupling $\xi > 0$ is mapped upon a downwards-oriented half-oval in the $\alpha - \beta$ plane with a top at $\alpha = \pi/4$. Its two asymptotes $\alpha = 0$ and $\alpha = \pi/2$ are reached in the limit $\beta \rightarrow -\infty$.

In the new graphical representation the robust, ξ -independent energy level $F = 0$ lies on the top of the half-oval while its decreasing and increasing neighbors are found displaced to the left and right, respectively, along the half-oval downwards. At the first few lowest $N = 2n+4$ the coordinates of these eigenvalues remain non-numerical,

$$\begin{aligned} F_0 = 0, \quad F_{\pm} &= \pm \sqrt{2 - \xi^2}, \quad n = 0, \\ F_0 = 0, \quad F_{\pm, \pm} &= \pm \sqrt{2 - \xi^2 \pm \sqrt{1 - 4\xi^2}}, \quad n = 1 \end{aligned}$$

etc. The closed form of these definitions enables us to determine the closed form of the respective critical values at which the spectrum ceases to be real,

$$\begin{aligned} Z_{crit}(4) &= 4\sqrt{2} \approx 5.66, \quad n = 0, \\ Z_{crit}(6) &= 9/2 = 4.50, \quad n = 1, \end{aligned}$$

followed by the numerically calculated $Z_{crit}(8) \approx 4.463$ (at $n = 2$), $Z_{crit}(10) \approx 4.461$ (at $n = 3$), $Z_{crit}(12) \approx 4.463$ (at $n = 4$) etc. These results do not contradict the expected $n \rightarrow \infty$ limit $Z_{crit}(\infty) \approx 4.475$ derived in ref. [12].

3.1 A real-matrix re-arrangement of eq. (8)

We may split eq. (8) in its real and imaginary parts and ‘glue’ them together in the following very natural pentadiagonal or, if you wish, block-tridiagonal eigenvalue

problem

$$\left(\begin{array}{cc|cc|c|c} -F & -\xi & -1 & 0 & & \\ \xi & -F & 0 & -1 & & \\ \hline -1 & 0 & -F & -\xi & \ddots & \\ 0 & -1 & \xi & -F & & \ddots \\ \hline & & \ddots & & \ddots & \ddots & -1 & 0 \\ & & & & \ddots & \ddots & 0 & -1 \\ \hline & & & & -1 & 0 & -F & -\xi & -1 \\ & & & & 0 & -1 & \xi & -F & 0 \\ \hline & & & & & & -2 & 0 & -F \end{array} \right) \left(\begin{array}{c} a_0 \\ b_0 \\ \hline a_1 \\ b_1 \\ \hline \vdots \\ \vdots \\ \hline a_n \\ b_n \\ \hline \gamma \end{array} \right) = 0. \quad (20)$$

This equation may be re-written in the partitioned-matrix notation,

$$\left(\begin{array}{cccc} \mathbf{X} & -\mathbf{1} & & \\ -\mathbf{1} & \mathbf{X} & \ddots & \\ & \ddots & \ddots & -\mathbf{1} \\ & & -\mathbf{1} & \mathbf{X} & \vec{\mathbf{d}} \\ & & & 2\vec{\mathbf{d}}^T & -F \end{array} \right) \left(\begin{array}{c} \vec{\mathbf{c}}_0 \\ \vec{\mathbf{c}}_1 \\ \vdots \\ \vec{\mathbf{c}}_n \\ \gamma \end{array} \right) = 0. \quad (21)$$

The obvious boldface two-by-two matrix elements degenerate, in an ‘odd’ anomalous last row and column, to an auxiliary vector $\vec{\mathbf{d}}^T = (1, 0)$.

A few comments are due. Firstly, all our wave-function components are now re-interpreted as proportional to the classical Tshebyshev polynomials U_k with a two-by-two matrix argument X . Equation (21) leads to an alternative formula for the eigenvectors,

$$\vec{\mathbf{c}}_{\mathbf{k}} = U_k \left(\frac{1}{2} \mathbf{X} \right) \vec{\mathbf{c}}_0, \quad \mathbf{X} = \begin{pmatrix} -F & -\xi \\ \xi & -F \end{pmatrix}, \quad \mathbf{k} = 0, 1, \dots, n+1. \quad (22)$$

Secondly, once we introduce a *complex* angle α we may parametrize $F = -\varrho \cos \alpha$ and $\xi = \varrho \sin \alpha$ using an optional, redundant parameter ϱ . A peculiar feature of our matrices X is that their powers remain elementary in this representation,

$$\mathbf{X}^m = \varrho^m \begin{pmatrix} \cos m\alpha & -\sin m\alpha \\ \sin m\alpha & \cos m\alpha \end{pmatrix}.$$

This means that all the formulae containing polynomials (22) remain amazingly transparent.

The existence of the explicit solutions (22) reduces eq. (21) to the two secular-equation constraints imposed upon the vector $\vec{\mathfrak{C}}_{n+1}$. Of course, they are equivalent to our complex matching conditions as mentioned above.

4 Solutions at the odd $N = 2n + 3$

Weigert [9] did not notice that a “one-step easier” discretization of eq. (4) emerges at the odd $N = 2n + 3$, with $2N + 2$ energy roots at $n \geq 0$. An alternative to eq. (8) then reads, in the same notation,

$$\left(\begin{array}{ccc|ccc} i\xi - F & -1 & & & & \\ & -1 & i\xi - F & \ddots & & \\ & & \ddots & \ddots & -1 & \\ & & & -1 & i\xi - F & -1 \\ \hline & & & -1 & & \\ & & & & -i\xi - F & \ddots \\ & & & & \ddots & \ddots & -1 \\ & & & & & -1 & -i\xi - F \end{array} \right) \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \vdots \\ \alpha_n \\ \alpha_n^* \\ \vdots \\ \alpha_0^* \end{pmatrix} = 0. \quad (23)$$

Definition (9) of the eigenvectors remains unchanged but the matching condition is just one,

$$\gamma = U_{n+1} \left(\frac{-F + i\xi}{2} \right) (a + ib) = U_n \left(\frac{-F - i\xi}{2} \right) (a - ib). \quad (24)$$

The ratio between this equation and its Hermitian conjugate eliminates all the normalization ambiguities and leads to the odd- N counterpart of eq. (16),

$$U_n \left(\frac{-F + i\xi}{2} \right) U_n \left(\frac{-F - i\xi}{2} \right) = U_{n+1} \left(\frac{-F + i\xi}{2} \right) U_{n+1} \left(\frac{-F - i\xi}{2} \right). \quad (25)$$

This secular equation is our final result. As an implicit definition of the $N = 2n + 3$ energy levels $F = F(\xi)$ it possesses the compact non-numerical solutions at the first two values of n again,

$$\begin{aligned} F_{\pm} &= \pm \sqrt{1 - \xi^2}, & n = 0, \\ F_{\pm, \pm} &= \pm \frac{1}{2} \sqrt{6 - 4\xi^2 \pm 2\sqrt{5 - 16\xi^2}}, & n = 1. \end{aligned}$$

The respective elementary expressions for the critical constants

$$\begin{aligned} Z_{crit}(3) &= \frac{9}{4} = 2.25, & n = 0, \\ Z_{crit}(5) &= \frac{25\sqrt{5}}{16} \approx 3.49, & n = 1 \end{aligned}$$

are followed by the complex Cardano representation of the real $Z_{crit}(7) \approx 3.946$ at $n = 2$. At $n > 2$ one switches to a purely numerical algorithm giving $Z_{crit}(9) \approx 4.148$ at $n = 3$ etc. In comparison with the parallel results sampled in section 3 at even N we notice a slowdown of the numerical convergence towards the $n \rightarrow \infty$ limit.

Acknowledgment

Proportionally supported by NPI, IRP AV0Z10480505, and by GA AS, contract No. A 1048302.

References

- [1] C. M. Bender and S. Boettcher, Phys. Rev. Lett. 80 (1998) 5243;
C. M. Bender, S. Boettcher and P. N. Meisinger, J. Math. Phys. 40 (1999) 2201.
- [2] F. G. Scholtz, H. B. Geyer and F. J. W. Hahne, Ann. Phys. (NY) 213 (1992) 74.
- [3] M. Znojil, Phys. Lett. A 259 (1999) 220;
M. Znojil, math-ph/0104012.
- [4] C. M. Bender, D. C. Brody and H. F. Jones, Phys. Rev. Lett. 89 (2002) 270401;
C. M. Bender, Czech. J. Phys. 54 (2004) 1027.
- [5] A. Mostafazadeh, J. Math. Phys. 43 (2002) 205 and 2814.
- [6] R. Kretschmer and L. Szymanowski, quant-ph/0105054;
P. Dorey, C. Dunning and R. Tateo, J. Phys. A: Math. Gen. 34 (2001) 5679;
K. C. Shin, Commun. Math. Phys. 229 (2002) 543;
F. Kleefeld, in “Hadron Physics, Effective Theories of Low Energy QCD”, AIP Conf. Proc. 660 (2003) 325;
V. Jakubský, Czech. J. Phys. 54 (2004) 67;
G. Sclarici, Czech. J. Phys. 54 (2004) 119;
Q. Wang, Czech. J. Phys. 54 (2004) 143;
A. Blasi, G. Sclarici and L. Solombrino, Czech. J. Phys. 54 (2004) 1055;
H. B. Geyer, F. G. Scholz and I. Snyman, Czech. J. Phys. 54 (2004) 1069;
W. D. Heiss, Czech. J. Phys. 54 (2004) 1091;
E. Caliceti, Czech. J. Phys. 55 (2005) 1077;
U. Günther and F. Stefani, Czech. J. Phys. 55 (2005) 1099;
A. Mostafazadeh, Czech. J. Phys. 55 (2005) 1157.

- [7] H. Langer and Ch. Tretter, Czech. J. Phys. 54 (2004) 1113.
- [8] M. Znojil, Phys. Lett. A. 285 (2001) 7;
M. Znojil, J. Math. Phys. 45 (2004) 4418.
- [9] S. Weigert, Czech. J. Phys. 55 (2005) 1183.
- [10] B. Bagchi, S. Mallik and C. Quesne, Mod. Phys. Lett. A17 (2002) 1651;
M. Znojil, J. Phys. A: Math. Gen. 36 (2003) 7825;
A. Mostafazadeh and A. Batal, J. Phys. A: Math. Gen. 37 (2004) 11645;
V. Jakubský and M. Znojil, Czech. J. Phys. 54 (2004) 1101;
M. Znojil, J. Math. Phys. 46 (2005) 062109;
H. Bíla, V. Jakubský, M. Znojil, B. Bagchi, S. Mallik and C. Quesne, Czech. J. Phys. 55 (2005) 1075.
- [11] G. Szegő, Orthogonal Polynomials, AMS, Providence, 1991.
- [12] M. Znojil and G. Lévai, Mod. Phys. Lett. A 16 (2001) 2273.